Author Index

AALBERSE R.C. *see* VAN NEERVEN R.J.J.
ALFANI E. *see* PAROLI M.
ALONSO L. *see* VICENTE A.
ANDERSON J.L. *see* CARLQUIST J.F.
ANDONEGUI G.A. *see* TREVANI A.S.
ANTEL J. *see* ULVESTAD E.
ANWAR A.R.E., WALSH G.M., CROMWELL O., KAY A.B. & WARDLAW A.J. Adhesion to fibronectin primes eosinophils via $\alpha_4\beta_1$ (VLA-4), 222
APPELBERG R., CASTRO A.G., PEDROSA J. & MINÓPRIO P. Role of interleukin-6 in the induction of protective T cells during mycobacterial infections in mice, 361
ARIZONO N. *see* UCHIKAWA R.
ARNOLD R., HUMBERT B., WERCHAU H., GALLATI H. & KÖNIG W. Interleukin-8, interleukin-6 and soluble tumour necrosis factor receptor type I release from a human pulmonary epithelial cell line (A549) exposed to respiratory syncytial virus, 126
ARNOLD R., WERNER F., HUMBERT B., WERCHAU H. & KÖNIG W. Effect of respiratory syncytial virus—antibody complexes on cytokine (IL-8, IL-6, TNF-α) release and respiratory burst in human granulocytes, 184
ASHERSON G.L. *see* DIELI F.
AUBERGER P. *see* MARI B.
AUBRY J.-P. *see* BACON K.B.

BACON K.B., FORES-ROMO L., AUBRY J.-P., WELLS T.N.C. & POWER C.A. Interleukin-8 and RANTES induce the adhesion of the human basophilic cell line KU-812 to human endothelial cell monolayers, 473
BAILEY M., HALL L., BLAND P.W. & STOKES C.R. Production of cytokines by lymphocytes from spleen, mesenteric lymph node and intestinal lamina propria of pigs, 577
BALABAUD C. *see* GARCIA-BARCINA M.
BANKS E.M.S. *see* HOLLIDAY M.R.
BARANJI K. *see* BARANYI L.
BARANYI L., BARANJI K., TAKIZAWA H., OKADA N. & OKADA H. Cell-surface bound complement regulatory activity is necessary for the *in vivo* survival of KDH-8 rat hepatoma, 522
BARNABA V. *see* PAROLI M.
BARRERA L.F., KRAMNIK I., SKAMENE E. & RADZIOCH D. Nitrite production by macrophages derived from BCG-resistant and -susceptible congenic mouse strains in response to IFN-γ and infection with BCG, 457
BAUGHN R.E. *see* WICHER V.
BELHACENE N. *see* MARI B.
BELLABY T. *see* ROBINSON K.
BELLONE M. *see* KARACHUNSKI P.I.
BENIGNI F. *see* MENGOZZI M.
VAN DEN BERG W.B. *see* JACOBS M.J.M.
BETJES M. *see* O'DOHERTY U.
BHARDWAJ N. *see* O'DOHERTY U.
BICKLE Q.D. *see* SMITH D.A.
BIDAURRAZAGA I. *see* GARCIA-BARCINA M.
BIOULAC-SAGE *see* GARCIA-BARCINA M.
BJØRGE L. *see* VEDELER C.
BJÖRK P. *see* CLINCHY B.
BJÖRKLUND M. *see* SUNDSTEDT A.
BLACKWELL J.M. *see* FORMICA S.
BLANCHARD-CHANNELL M., MOORE P.F. & STOTT J.L. Characterization of monoclonal antibodies specific for equine homologues of CD3 and CD5, 548
BLAND P.W. *see* BAILEY M.
BØ L. *see* ULVESTAD E.
BOOTS A.H.M., WIMMERS-BERTENS A.J.M.M. & RIJNDERS A.W.M. Antigen-presenting capacity of rheumatoid synovial fibroblasts, 268
BOUSQUET J. *see* CAMPBELL A.M.
BOYD C.A.R. *see* CRAWFORD D.H.
BREITTMAYER J.-P. *see* MARI B.
BROSSARD M. *see* MBOW M.L.
BUJDOSO R. *see* EVANS C.W.

CAI Z. *see* HU Z.-Q.
CALDER P.C. *see* YAQOOB P.
CALZETTI F. *see* SERRA M.C.
CAMPBELL A.K. *see* DAVIES E.V.
CAMPBELL A.M., VIGNOLA A.M., CHANEZ P., GODARD Ph. & BOUSQUET J. Low-affinity receptor for IgE on human bronchial epithelial cells in asthma, 506
CARLONI G. *see* PAROLI M.
CARLQUIST J.F., SHELBY J., HAMMOND E.H., GREENWOOD J.H. & ANDERSON J.L. Histoincompatibility-associated differences in the phenotype of murine cardiac allograft infiltrating T cells, 149
CASSATELLA M.A. *see* SERRA M.C.
CASTRO A.G. *see* APPELBERG R.
CESKA M. *see* SERRA M.C.
CHANDLER P. *see* O'REILLY L.A.
CHANEZ P. *see* CAMPBELL A.M.
CHEN S. *see* CRAWFORD D.H.
CHENG Y.F. *see* DICK A.D.
CIRCOLO A. *see* KULICS J.
CLARK B.D. *see* MENGOZZI M.
CLINCHY B., BJÖRK P., PAULIE S. & MÖLLER G. Interleukin-10 inhibits motility in murine and human B lymphocytes, 376
COLE F.S. *see* STRUNK R.C.
COLEMAN J.W. *see* HOLLIDAY M.R.
COLTEN H.R. *see* KULICS J.
CONTI G. *see* VEDELER C.
CONTI-TRONCONI B.M. *see* KARACHUNSKI P.I.
COOKE A. *see* O'REILLY L.A.
COTTERILL T. *see* STEWART A.G.
CRAWFORD D.H., CHEN S. & BOYD C.A.R. Cationic amino acid transport in human T lymphocytes is markedly increased in the CD45RA, $CD8^+$ population after activation, 357
see also LAM K.M.C.
CROMWELL O. *see* ANWAR A.R.E.
CZARNNETZKI B.M. *see* WEBER S.

DALUM I. *see* MOURITSEN S.
DANIELS R.H., ELMORE M.A., HILL M.E., SHIMIZU Y., LACKIE J.M. & FINNEN M.J. Priming of the oxidative burst in human neutrophils by physiological agonists or cytochalasin B results from the recruitment of previously non-responsive cells, 465
DAVIES E.V., CAMPBELL A.K. & HALLETT M.B. Ca^{2+} oscillations in neutrophils triggered by immune complexes result from Ca^{2+} influx, 57
DEARMAN R.J. *see* HOLLIDAY M.R.
DECKERT M. *see* MARI B.
DEL CARPIO C. *see* DIELI F.
DENIS M. Neutralization of transforming growth factor-β_1 in a mouse model of immune-induced lung fibrosis, 584
DICK A.D., CHENG Y.F., LIVERSIDGE J. & FORRESTER J.V. Intranasal administration of retinal antigens suppresses retinal antigen-induced experimental autoimmune uveoretinitis, 625
DIELI F., ASHERSON G.L., SIRECI G., TANTILLO G., DEL CARPIO C. & SALERNO A. Dominant Vβ8 gene usage in

response to TNP: failure to use other Vβ chains following removal of Vβ8$^+$ T cells by monoclonal antibody *in vivo*, 99
DIPPEL E. *see* WEBER S.
DOHLSTEN M. *see* SUNDSTEDT A.
DOUGLAS A.J. *see* POLLOCK J.M.
DOUGLAS J.T. *see* ELZER P.H.
DOUGLAS M.W., KESSON A.M. & KING N.J.C. CTL recognition of West Nile virus-infected fibroblasts is cell cycle dependent and is associated with virus-induced increases in class I MHC antigen expression, 561
DOWNING J.E.G. Multiple nitric oxide synthase systems in adult rat thymus revealed using NADPH diaphorase histochemistry, 659
DURHAM S.R. *see* MASUYAMA K.
see also YING S.

EHNHOLM C. *see* VÄKEVÄ A.
ELMORE M.A. *see* DANIELS R.H.
ELSNER H. *see* MOURITSEN S.
ELZER P.H., JACOBSON R.H., JONES S.M., NIELSEN K.H., DOUGLAS J.T. & WINTER A.J. Antibody-mediated protection against *Brucella abortus* in BALB/c mice at successive periods after infection: variation between virulent strain 2308 and attenuated vaccine strain 19, 651
ENGEL A.M. *see* MOURITSEN S.
EVANS C.W., LUND B.T., McCONNELL I. & BUJDOSO R. Antigen recognition and activation of ovine γδ cells, 229

FAGGIONI R. *see* MENGOZZI M.
FALINI B. *see* PULFORD K.
FALK K., RÖTZSCHKE O., STEVANOVIĆ S., GNAU V., SPARBIER K., JUNG G., RAMMENSEE H.-G. & WALDEN P. Analysis of a naturally occurring HLA class I-restricted viral epitope, 337
FANTUZZI G. *see* MENGOZZI M.
FAURE P., MARUSSIG M., GOOSSENS P., MILTGEN F. & MAZIER D. Characterization of liver lymphomyeloid cells in mice infected with *Plasmodium yoelii* sporozoites, 645
FINNEN M.J. *see* DANIELS R.H.
FLEISCHER J.A. *see* STRUNK R.C.
FORES-ROMO L. *see* BACON K.B.
FORMICA S., ROACH T.I.A. & BLACKWELL J.M. Interaction with extracellular matrix proteins influences *Lsh/Ity/Bcg* (candidate *Nramp*) gene regulation of macrophage priming/activation for tumour necrosis factor-α and nitrite release, 42
FORRESTER J.V. *see* DICK A.D.
FRANCO A. *see* PAROLI M.
FRENCH L.E., SAPPINO A.-P., TSCHOPP J. & SCHIFFERLI J.A. Clusterin gene expression in the rat thymus is not modulated by dexamethasone treatment, 328
FRIEDL P., NOBLE P.B., SHIELDS E.D. & ZÄNKER K.S. Locomotor phenotypes of unstimulated CD45RA^{high} and CD45RO^{high} CD4$^+$ and CD8$^+$ lymphocytes in three-dimensional collagen lattices, 617
FRISCHHOLZ S., RÖLLINGHOFF M. & MOLL H. Cutaneous leishmaniasis: co-ordinate expression of granzyme A and lymphokines by CD4$^+$ T cells from susceptible mice, 255
FUCHSBERGER N. *see* KUBEŠ M.
VAN FURTH R. *see* GEERTSMA M.F.

GABRILOVICH D.I., WOODS G.M., PATTERSON S., HARVEY J.J. & KNIGHT S.C. Retrovirus-induced immunosuppression via blocking of dendritic cell migration and down-regulation of adhesion molecules, 82
GALANOS C. *see* HONARVAR N.
GALLATI H. *see* ARNOLD R.
GARCIA-BARCINA M., WINNOCK M., BIDAURRAZAGA I., HUET S., BIOULAC-SAGE & BALABAUD C. Detection of cell-adhesion molecules on human liver-associated lymphocytes, 95
GARSIDE P., REID S., STEEL M. & MOWAT A. McI. Differential cytokine production associated with distinct phases of murine graft-versus-host reaction, 211
GEERTSMA M.F., TEEUW W.L., NIBBERING P.H. & VAN FURTH R. Pulmonary surfactant inhibits activation of human monocytes by recombinant interferon-γ, 450
GEFFNER J.R. *see* TREVANI A.S.
GEZELTER S. *see* O'DOHERTY U.
GHEZZI P. *see* MENGOZZI M.
GIBBONS A.E., PRICE P. & SHELLAM G.R. Bone marrow atrophy induced by murine cytomegalovirus infection, 410
GLASS E.J. & MILLER P. Induction of effective cross-reactive immunity by FMDV peptides is critically dependent upon specific MHC–peptide–T cell interactions, 1
GNAU V. *see* FALK K.
GODARD Ph. *see* CAMPBELL A.M.
GOLDMAN M. *see* MENGOZZI M.
GÓMEZ DE MORAL M. *see* VICENTE A.
GOOSSENS P. *see* FAURE P.
GÖTZE O. *see* OPPERMANN M.
GRAHAM C.M. *see* SMITH C.A.
GREENWOOD J.H. *see* CARLQUIST J.F.
GRZYWACZ M. *see* LAM K.M.C.
GUERIN S. *see* MARI B.
GUSHIMA H. *see* SUGITA Y.

HAAK-FRENDSCHO M., ROBBINS K., LYON R., SHIELDS R., HOOLEY J., SCHOENHOFF M. & JARDIEU P. Administration of an anti-IgE antibody inhibits CD23 expression and IgE production *in vivo*, 306
HALL L. *see* BAILEY M.
HALLETT M.B. *see* DAVIES E.V.
HAMAD M. & KLEIN J.R. Phenotypic and functional heterogeneity of murine intestinal intraepithelial lymphocytes defined by cell density: implications for route of differentiation and responsiveness to proliferation induction, 611
HAMID Q. *see* MASUYAMA K.
HAMID Q.A. *see* YING S.
HAMMOND E.H. *see* CARLQUIST J.F.
HAMPRECHT K. & STEINMASSL M. Low-dose human cytomegalovirus infection of human fibroblast cultures induces lymphokine-activated killer cell resistance: interferon-β-mediated target cell protection does not correlate with up-regulation of HLA class I surface molecules, 171
HARRIS T. *see* STEWART A.G.
HARVEY J.J. *see* GABRILOVICH D.I.
HE X. *see* KIRSCHMANN D.A.
HEALEY D. *see* O'REILLY L.A.
HEDLUND G. *see* SUNDSTEDT A.
HERNANDEZ-PANDO R. & ROOK G.A.W. The role of TNF-α in T-cell-mediated inflammation depends on the Th1/Th2 cytokine balance, 591
HETLAND G. & WIKER H.G. Antigen 85C on *Mycobacterium bovis*, BCG and *M. tuberculosis* promotes monocyte-CR3-mediated uptake of microbeads coated with mycobacterial products, 445
HILL M.E. *see* DANIELS R.H.
HIROMATSU K. *see* TAKADA H.
VAN DEN HOEK A.E.M. *see* JACOBS M.J.M.
HÖIDÉN I. *see* SUNDSTEDT A.
HOLLIDAY M.R., BANKS E.M.S., DEARMAN R.J., KIMBER I. & COLEMAN J.W. Interactions of IFN-γ with IL-3 and IL-4 in the regulation of serotonin and arachidonate release from mouse peritoneal mast cells, 70
HONARVAR N., SCHAIBLE U.E., GALANOS C., WALLICH R. & SIMON M.M. A 14,000 MW lipoprotein and a glycolipid-like structure of *Borrelia burgdorferi* induce proliferation and immunoglobulin production in mouse B cells at high frequencies, 389
HOOLEY J. *see* HAAK-FRENDSCHO M.
HU Z.-Q., YAMAZAKI T., CAI Z., YOSHIDA T. & SHIMAMURA T. Mast cells display natural suppressor activity partially by releasing transforming growth factor-β, 482
HUET S. *see* GARCIA-BARCINA M.
HUMBERT B. *see* ARNOLD R.
HYODOH F. *see* UEKI A.

IANARO A., O'DONNELL C.A., DI ROSA M. & LIEW F.Y. A

nitric oxide synthase inhibitor reduces inflammation, down-regulates inflammatory cytokines and enhances interleukin-10 production in carrageenin-induced oedema in mice, 370
IKEDA H. *see* TAGUCHI O.
ISTURIZ M.A. *see* TREVANI A.S.
ITO K. *see* SUGITA Y.
IWAMOTO I. *see* KITA Y.

JACOBS M.J.M., VAN DEN HOEK A.E.M., VAN DE PUTTE L.B.A. & VAN DEN BERG W.B. Anergy of antigen-specific T lymphocytes is a potent mechanism of intravenously induced tolerance, 294
JACOBSON M.R. *see* MASUYAMA K.
see also YING S.
JACOBSON R.H. *see* ELZER P.H.
JANSEN H.M. *see* VAN NEERVEN R.J.J.
JARDIEU P. *see* HAAK-FRENDSCHO M.
JAUHIAINEN M. *see* VÄKEVÄ A.
JONES J. & MORGAN B.P. Comparative susceptibility of peripheral blood leucocytes and related cell lines to killing by T-cell perforin, 555
JONES M. *see* PULFORD K.
JONES S.M. *see* ELZER P.H.
JUNG G. *see* FALK K.

KALLAND T. *see* SUNDSTEDT A.
KAPLAN A. *see* QUDDUS J.
KAPSENBERG M.L. *see* VAN NEERVEN R.J.J.
KARACHUNSKI P.I., OSTLIE N., CONTI-TRONCONI B.M. & BELLONE M. Residues with the α subunit sequence 304–322 of muscle acetylcholine receptor forming autoimmune $CD4^+$ epitopes in BALB/c mice, 22
KATZ Y. *see* STRUNK R.C.
KAWAKAMI Y. *see* UEKI A.
KAY A.B. *see* ANWAR A.R.E.
see also MASUYAMA K., YING S.
KAYE P.M. *see* SMITH D.A.
KESSON A.M. *see* DOUGLAS M.W.
KEZUKA T. *see* TAGUCHI O.
KIMBER I. *see* HOLLIDAY M.R.
KING N.J.C. *see* DOUGLAS M.W.
KINUGAWA K. *see* UEKI A.
KIRSCHMANN D.A., HE X. & MURASKO D.M. Inhibition of macrophage-induced, antigen-specific T-cell proliferation by poly I:C role of suppressor macrophages, 238
KITA Y., SUMIDA T., IWAMOTO I., YOSHIDA S. & KOIKE T. V gene analysis of anti-cardiolipin antibodies from (NZW × BSXB) F_1 mice, 494
KLEIN J.R. *see* HAMAD M.
KNIGHT S.C. *see* GABRILOVICH D.I.
KOIKE T. *see* KITA Y.
KOLLIAS G. *see* TAVERNE J.
KÖNIG W. *see* ARNOLD R.
KONTANI K. *see* TAGUCHI O.
KRAMNIK I. *see* BARRERA L.F.
KUBEŠ M., FUCHSBERGER N., LABUDA M., ŽUFFOVÁ E. & NUTTALL P.A. Salivary gland extracts of partially fed *Dermacentor reticulatus* ticks decrease natural killer cell activity *in vitro*, 113
KULICS J., CIRCOLO A., STRUNK R.C. & COLTEN H.R. Regulation of synthesis of complement protein C4 in human fibroblasts: cell- and gene-specific effects of cytokines and lipopolysaccharide, 509
KURODA A. *see* UCHIKAWA R.

LABUDA M. *see* KUBEŠ M.
LACKIE J.M. *see* DANIELS R.H.
LAM K.M.C., WHITTLE H., GRZYWACZ M. & CRAWFORD D.H. Epstein–Barr virus-carrying B cells are large, surface IgM, IgD-bearing cells in normal individuals and acute malaria patients, 383
LEHTO T. *see* VÄKEVÄ A.
LIEW F.Y. *see* IANARO A.
LINKE A.T. & MALE D.K. Strain-specific variation in constitutive and inducible expression of MHC class II, class I and ICAM-1 on rat cerebral endothelium, 88
LIU Z.-Z. *see* SATOH S.
LIVERSIDGE J. *see* DICK A.D.
LOPEZ BERNAL A. *see* MALHOTRA R.
LOWHAGEN O. *see* MASUYAMA K.
see also YING S.
LOWRIE D.B. *see* SILVA C.L.
LUND B.T. *see* EVANS C.W.
LUND T. *see* O'REILLY L.A.
LYON R. *see* HAAK-FRENDSCHO M.

MacKIE D.P. *see* POLLOCK J.M.
McCONNELL I. *see* EVANS C.W.
MALE D.K. *see* LINKE A.T.
MALHOTRA R., WILLIS A.C., LOPEZ BERNAL A., THIEL S. & SIM R.B. Mannan-binding protein levels in human amniotic fluid during gestation and its interaction with collectin receptor from amnion cells, 439
MARCHANT A. *see* MENGOZZI M.
MARI B., BREITTMAYER J.-P., GUERIN S., BELHACENE N., PEYRON J.-F., DECKERT M., ROSSI B. & AUBERGER P. High levels of functional endopeptidase 24.11 (CD10) activity on human thymocytes: preferential expression on immature subsets, 433
MARUSSIG M. *see* FAURE P.
MASON D.Y. *see* PULFORD K.
MASUHO Y. *see* SUGITA Y.
MASUYAMA K., JACOBSON M.R., RAK S., MENG Q., SUDDERICK R.M., KAY A.B., LOWHAGEN O., HAMID Q. & DURHAM S.R. Topical glucocorticosteroid (fluticasone propionate) inhibits cells expressing cytokine mRNA for interleukin-4 in the nasal mucosa in allergen-induced rhinitis, 192
see also YING S.
MATRE R. *see* VEDELER C.
MATSUDA S. *see* UCHIKAWA R.
MATSUZAKI G. *see* TAKADA H.
MAYNE K.M. *see* PULFORD K.
MAZIER D. *see* FAURE P.
MBOW M.L., RUTTI B. & BROSSARD M. Infiltration of $CD4^+$ $CD8^+$ T cells, and expression of ICAM-1, Ia antigens, IL-1α and TNF-α in the skin lesion of BALB/c mice undergoing repeated infestations with nymphal *Ixodes ricinus* ticks, 596
MENG Q. *see* MASUYAMA K.
MENGOZZI M., FANTUZZI G., FAGGIONI R., MARCHANT A., GOLDMAN M., ORENCOLE S., CLARK B.D., SIRONI M., BENIGNI F. & GHEZZI F. Chlorpromazine specifically inhibits peripheral and brain TNF production, and up-regulates IL-10 production, in mice, 207
MERI S. *see* VÄKEVÄ A.
MERINO J. *see* RAMOS A.
MERINO R. *see* RAMOS A.
MICKLEM K.J. *see* PULFORD K.
MILLER P. *see* GLASS E.J.
MILTGEN F. *see* FAURE P.
MINÓPRIO P. *see* APPELBERG R.
MOLL H. *see* FRISCHHOLZ S.
MÖLLER G. *see* CLINCHY B.
MOORE P.F. *see* BLANCHARD-CHANNELL M.
MORGAN B.P. *see* JONES J.
MØRK S. *see* ULVESTAD E.
see also VEDELER C.
MORRISON H. *see* PULFORD K.
MOURITSEN S., DALUM I., ENGEL A.M., SVANE I.-M., SVENDSEN I., WERDELIN O. & ELSNER H. MHC class II-bound self-peptides can be effectively separated by isoelectric focusing and bind optimally to their MHC class II restriction elements around pH 5·0, 529
MOWAT A. McI. *see* GARSIDE P.
MURASKO D.M. *see* KIRSCHMANN D.A.

VAN NEERVEN R.J.J., VAN DE POL M.M., WIERENGA E.A.,

AALBERSE R.C., JANSEN H.M. & KAPSENBERG M.L. Peptide specificity and HLA restriction do not dictate lymphokine production by allergen-specific T-lymphocyte clones, 351
NEILL S.D. *see* POLLOCK J.M.
NEWMAN I. *see* WILKINSON P.C.
NEWSHOLME E.A. *see* YAQOOB P.
NIBBERING P.H. *see* GEERTSMA M.F.
NIELSEN K.H. *see* ELZER P.H.
NIKOLOVA E.B., TOMANA M. & RUSSELL M.W. The role of the carbohydrate chains in complement (C3) fixation by solid-phase-bound human IgA, 321
NOBLE P.B. *see* FRIEDL P.
NOMOTO K. *see* TAKADA H.
NORTON A. *see* PULFORD K.
NUSSENZWEIG V. *see* RÉNIA L.
NÜSSLER A.K. *see* SATOH S.
NUTTALL P.A. *see* KUBEŠ M.

OBEID O.E. & STEWARD M.W. The potential of immunization with synthetic peptides to overcome the immunosuppressive effect of maternal anti-measles virus antibodies in young mice, 16
O'DOHERTY U., PENG M., GEZELTER S., SWIGGARD W.J., BETJES M., BHARDWAJ N. & STEINMAN R.M. Human blood contains two subsets of dendritic cells, one immunologically mature and the other immature, 487
O'DONNELL C.A. *see* IANARO A.
OHSAWA G. *see* UEKI A.
OKADA H. *see* BARANYI L.
OKADA N. *see* BARANYI L.
OPPERMANN M. & GÖTZE O. Plasma clearance of the human C5a anaphylatoxin by binding to leucocyte C5a receptors, 516
O'REILLY L.A., HEALEY D., SIMPSON E., CHANDLER P., LUND T., RITTER M.A. & COOKE A. Studies on the thymus of non-obese diabetic (NOD) mice: effect of transgene expression, 275
ORENCOLE S. *see* MENGOZZI M.
OSTLIE N. *see* KARACHUNSKI P.I.

PAROLI M., CARLONI G., FRANCO A., DE PETRILLO E., ALFANI E., PERRONE A. & BARNABA V. Human hepatoma cells expressing MHC antigens display accessory cell function: dependence on LFA-1/ICAM-1 interaction, 215
PATTERSON S. *see* GABRILOVICH D.I.
PAULIE S. *see* CLINCHY B.
PEDROSA J. *see* APPELBERG R.
PENG M. *see* O'DOHERTY U.
PERRONE A. *see* PAROLI M.
DE PETRILLO E. *see* PAROLI M.
PEYRON J.-F. *see* MARI B.
PEZZELLA F. *see* PULFORD K.
PLAYFAIR J.H.L. *see* TAVERNE J.
VAN DE POL M.M. *see* VAN NEERVEN R.J.J.
POLLOCK J.M., DOUGLAS A.J., MacKIE D.P. & NEILL S.D. Identification of bovine T-cell epitopes for three *Mycobacterium bovis* antigens: MPB70, 19,000 MW and MPB57, 9
POWER C.A. *see* BACON K.B.
VAN DE PUTTE L.B.A. *see* JACOBS M.J.M.
PRICE P. *see* GIBBONS A.E.
PROBERT L. *see* TAVERNE J.
PULFORD K., MICKLEM K.J., JONES M., PEZZELLA F., MAYNE K.M., MORRISON H., THOMAS J., FALINI B., NORTON A. & MASON D.Y. A novel internal antigen which distinguishes germinal centre cells from other B-cell types, 154

QUDDUS J., KAPLAN A. & RICHARDSON B.C. Anti-CD11a prevents deletion of self-reative T cells in neonatal C57BR mice, 301

RADZIOCH D. *see* BARRERA L.F.
RAK S. *see* MASUYAMA K.
see also YING S.
RAKHMILEVICH A.L. Evidence for a significant role of $CD4^+$ T cells in adoptive immunity to *Listeria monocytogenes* in the liver, 249
RAMMENSEE H.-G. *see* FALK K.
RAMOS A., MERINO R. & MERINO J. Differences in non-MHC alloantigens promote tissue rejection but fail to mediate allogeneic co-operation and autoimmunity in mice neonatally injected with semi-allogeneic F_1 B cells, 287
REID S. *see* GARSIDE P.
RÉNIA L., RODRIGUES M.M. & NUSSENZWEIG V. Intrasplenic immunization with infected hepatocytes: a mouse model for studying protective immunity against malaria pre-erythrocytic stage, 164
RICHARDSON B.C. *see* QUDDUS J.
RIJNDERS A.W.M. *see* BOOTS A.H.M.
RITTER M.A. *see* O'REILLY L.A.
ROACH T.I.A. *see* FORMICA S.
ROBBINS K. *see* HAAK-FRENDSCHO M.
ROBINSON K., BELLABY T. & WAKELIN D. Vaccination against the nematode *Trichinella spiralis* in high- and low-responder mice. Effects of different adjuvants upon protective immunity and immune responsiveness, 261
RODRIGUES M.M. *see* RÉNIA L.
RÖLLINGHOFF M. *see* FRISCHHOLZ S.
ROOK G.A.W. *see* HERNANDEZ-PANDO R.
DI ROSA M. *see* IANARO A.
ROSSI B. *see* MARI B.
RÖTZSCHKE O. *see* FALK K.
RUBIO N. Demonstration of the presence of an interleukin-1 receptor on the surface of murine astrocytes and its regulation by cytokines and Theiler's virus, 178
RUH B. *see* WEBER S.
RUSSELL M.W. *see* NIKOLOVA E.B.
RUTTI B. *see* MBOW M.L.

SALERNO A. *see* DIELI F.
SAPPINO A.-P. *see* FRENCH L.E.
SATOH S., NÜSSLER A.K., LIU Z.-Z. & THOMSON A.W. Proinflammatory cytokines and endotoxin stimulate ICAM-1 gene expression and secretion by normal human hepatocytes, 571
SCHAIBLE U.E. *see* HONARVAR N.
SCHATNER M. *see* TREVANI A.S.
SCHIFFERLI J.A. *see* FRENCH L.E.
SCHOENHOFF M. *see* HAAK-FRENDSCHO M.
SEREBRINSKY G. *see* TREVANI A.S.
SERRA M.C., CALZETTI F., CESKA M. & CASSATELLA M.A. Effect of substance P on superoxide anion and IL-8 production by human PMNL, 63
SHEIKH N. *see* TAVERNE J.
SHELBY J. *see* CARLQUIST J.F.
SHELLAM G.R. *see* GIBBONS A.E.
SHIELDS E.D. *see* FRIEDL P.
SHIELDS R. *see* HAAK-FRENDSCHO M.
SHIMAMURA T. *see* HU Z.-Q.
SHIMIZU T. & STREILEIN J.W. Evidence that ultraviolet B radiation induces tolerance and impairs induction of contact hypersensitivity by different mechanisms, 140
SHIMIZU Y. *see* DANIELS R.H.
SHIOZUKA K. *see* SUGITA Y.
SILVA C.L. & LOWRIE D.B. A single mycobacterial protein (hsp 65) expressed by a transgenic antigen-presenting cell vaccinates mice against tuberculosis, 244
SIM R.B. *see* MALHOTRA R.
SIMON M.M. *see* HONARVAR N.
SIMPSON E. *see* O'REILLY L.A.
SIRECI G. *see* DIELI F.
SIRONI M. *see* MENGOZZI M.
SKAMENE E. *see* BARRERA L.F.
SMITH C.A., GRAHAM C.M. & THOMAS D.B. Immunodominance correlates with T-cell receptor ($\alpha\beta$) gene usage in the class II-restricted response to influenza haemagglutinin, 343
SMITH D.A., BICKLE Q.D. & KAYE P.M. Resistance induced by drug abbreviated *Schistosoma mansoni* infections: treatment with the drug Ro11-3128 leads to enhanced antigen presentation, 419
DE SOUZA J.B. *see* TAVERNE J.
SPARBIER K. *see* FALK K.
STEEL M. *see* GARSIDE P.

STEINMAN R.M. *see* O'DOHERTY U.
STEINMASSL M. *see* HAMPRECHT K.
STEVANOVIĆ S. *see* FALK K.
STEWARD M.W. *see* OBEID O.E.
STEWART A.G., COTTERILL T. & HARRIS T. Granulocyte and granulocyte–macrophage colony-stimulating factors exert differential effects on neutrophil platelet-activating factor generation and release, 51
STOKES C.R. *see* BAILEY M.
STOTT J.L. *see* BLANCHARD-CHANNELL M.
STREILEIN J.W. *see* SHIMIZU T.
STRUNK R.C., FLEISCHER J.A., KATZ Y. & COLE F.S. Developmentally regulated effects of lipopolysaccharide on biosynthesis of the third component of complement and factor B in human fibroblasts and monocytes, 314
see also KULICS J.
SUDDERICK R.M. *see* MASUYAMA K.
SUGITA Y., ITO K., SHIOZUKA K., SUZUKI H., GUSHIMA H., TOMITA M. & MASUHO Y. Recombinant soluble CD59 inhibits reactive haemolysis with complement, 34
SUMIDA T. *see* KITA Y.
SUNDSTEDT A., DOHLSTEN M., HEDLUND G., HÖIDÉN I., BJÖRKLUND M. & KALLAND T. Superantigens anergize cytokine production but not cytotoxicity *in vivo*, 117
SUZUKI H. *see* SUGITA Y.
SVANE I.-M. *see* MOURITSEN S.
SVENDSEN I. *see* MOURITSEN S.
SWIGGARD W.J. *see* O'DOHERTY U.

TAGUCHI O., KONTANI K., IKEDA H., KEZUKA T., TAKEUCHI M., TAKAHASHI T. & TAKAHASHI T. Tissue-specific suppressor T cells involved in self-tolerance are activated extrathymically by self-antigens, 365
TAKADA H., MATSUZAKI G., HIROMATSU K. & NOMOTO K. Analysis of the role of natural killer cells in *Listeria monocytogenes* infection: relation between natural killer cells and T-cell receptor $\gamma\delta$ T cells in the host defence mechanism at the early stage of infection, 106
TAKAHASHI T. *see* TAGUCHI O.
TAKEUCHI M. *see* TAGUCHI O.
TAKIZAWA H. *see* BARANYI L.
TANTILLO G. *see* DIELI F.
TAVERNE J., SHEIKH N., DE SOUZA J.B., PLAYFAIR J.H.L., PROBERT L. & KOLLIAS G. Anaemia and resistance to malaria in transgenic mice expressing human tumour necrosis factor, 397
TEEUW W.L. *see* GEERTSMA M.F.
THIEL S. *see* MALHOTRA R.
THOMAS D.B. *see* SMITH C.A.
THOMAS J. *see* PULFORD K.
THOMSON A.W. *see* SATOH S.
TOMANA M. *see* NIKOLOVA E.B.
TOMITA M. *see* SUGITA Y.
TRAPP B. *see* ULVESTAD E.
TREVANI A.S., ANDONEGUI G.A., ISTURIZ M.A., SCHATNER M., SEREBRINSKY G. & GEFFNER J.R. Effect of proteolytic enzymes on neutrophil FcγRII activity, 632
TSCHOPP J. *see* FRENCH L.E.

UCHIKAWA R., YAMADA M., MATSUDA S., KURODA A. & ARIZONO N. IgE antibody production is associated with suppressed interferon-γ levels in mesenteric lymph nodes of rats infected with the nematode *Nippostrongylus brasiliensis*, 427
UEKI A., YAMAGUCHI M., UEKI H., WATANABE Y., OHSAWA G., KINUGAWA K., KAWAKAMI Y. & HYODOH F. Polyclonal human T-cell activation by silicate *in vitro*, 332
UEKI H. *see* UEKI A.
ULVESTAD E., WILLIAMS K., BØ L., TRAPP B., ANTEL J. & MØRK S. HLA class II molecules (HLA-DR, -DP, -DQ) on cells in the human CNS studied *in situ* and *in vitro*, 535
see also VEDELER C.

VÄKEVÄ A., JAUHIAINEN M., EHNHOLM C., LEHTO T. & MERI S. High-density lipoproteins can act as carriers of glycophosphoinositol lipid-anchored CD59 in human plasma, 28
VARAS A. *see* VICENTE A.
VEDELER C., ULVESTAD E., BJØRGE L., CONTI G., WILLIAMS K., MØRK S. & MATRE R. The expression of CD59 in normal human nervous tissue, 542
VICENTE A., VARAS A., ALONSO L., GÓMEZ DE MORAL M. & ZAPATA A.G. Ontogeny of rat thymic dendritic cells, 75
VIGNOLA A.M. *see* CAMPBELL A.M.

WAKELIN D. *see* ROBINSON K.
WALDEN P. *see* FALK K.
WALLICH R. *see* HONARVAR N.
WALSH G.M. *see* ANWAR A.R.E.
WARDLAW A.J. *see* ANWAR A.R.E.
WATANABE Y. *see* UEKI A.
WEBER S., RUH B., DIPPEL E. & CZARNNETZKI B.M. Monoclonal antibodies to leucosialin (CD43) induce homotypic aggregation of the human mast cell line HMC-1: characterization of leucosialin on HMC-1 cells, 638
WELLS T.N.C. *see* BACON K.B.
WERCHAU H. *see* ARNOLD R.
WERDELIN O. *see* MOURITSEN S.
WERNER F. *see* ARNOLD R.
WHITTLE H. *see* LAM K.M.C.
WICHER K. *see* WICHER V.
WICHER V., BAUGHN R.E. & WICHER K. Congenital and neonatal syphilis in guinea-pigs show a different pattern of immune response, 404
WIERENGA E.A. *see* VAN NEERVEN R.J.J.
WIKER H.G. *see* HETLAND G.
WILKINSON P.C. & NEWMAN I. Chemoattractant activity of IL-2 for human lymphocytes: a requirement for the IL-2 receptor β-chain, 134
WILLIAMS K. *see* ULVESTAD E.
see also VEDELER C.
WILLIS A.C. *see* MALHOTRA R.
WIMMERS-BERTENS A.J.M.M. *see* BOOTS A.H.M.
WINNOCK M. *see* GARCIA-BARCINA M.
WINTER A.J. *see* ELZER P.H.
WOODS G.M. *see* GABRILOVICH D.I.

YAMADA M. *see* UCHIKAWA R.
YAMAGUCHI M. *see* UEKI A.
YAMAZAKI T. *see* HU Z.-Q.
YAQOOB P., NEWSHOLME E.A. & CALDER P.C. The effect of dietary lipid manipulation on rat lymphocyte subsets and proliferation, 603
YING S., DURHAM S.R., JACOBSON M.R., RAK S., MASUYAMA K., LOWHAGEN O., KAY A.B. & HAMID Q.A. T lymphocytes and mast cells express messenger RNA for interleukin-4 in the nasal mucosa in allergen-induced rhinitis, 200
YOSHIDA S. *see* KITA Y.
YOSHIDA T. *see* HU Z.-Q.

ZÄNKER K.S. *see* FRIEDL P.
ZAPATA A.G. *see* VICENTE A.
ŽUFFOVÁ E. *see* KUBEŠ M.

Subject Index

Adhesion molecules:
  detection on human liver-associated lymphocytes, 95
Adjuvants:
  effects on protective immunity and immune responses to *Trichinella*, 261
Allergen-specific T-cell clones:
  lymphokine production by, 351
Alloantigens, non-major histocompatibility complex, role in tissue rejection:
  allogeneic co-operation and autoimmunity, 287
Amino acid transport, cationic:
  in human T cells, 357
Antigen presentation, enhanced:
  following treatment of *Schistosoma mansoni* with the drug Ro11-3128, 419
Antigen-presenting capacity:
  of rheumatoid synovial fibroblasts, 268
Autoimmune CD4 epitope:
  in the α subunits of acetylcholine receptor, 22

Borrelia burgdorferi:
  B-cell mitogens, 389
Brucella, protection against:
  by antibodies, 651

C3 and factor B biosynthesis:
  effects of lipopolysaccharide, 314
C5a:
  clearance from plasma, 516
Calcium oscillations:
  in neutrophils, 57
CD3 and CD5, equine homologues of:
  monoclonal antibodies to, 548
CD4 T cells, role in adoptive:
  immunity to *Listeria*, 249
CD10 expression:
  in human thymocytes, 433
CD23 expression:
  inhibition by anti-IgE, 306
CD59:
  expression in human nervous tissue, 542
CD59, glycophosphoinositol-anchored:
  high-density lipoproteins as carriers, 28
CD59, recombinant soluble:
  inhibition of reactive haemolysis, 34
Cerebral endothelium, rat:
  expressing major histocompatiblity complex ICAM-1, 88
Chlorpromazine:
  effect on tumour necrosis factor and interleukin-10 production, 207
Clusterin, gene expression:
  and dexamethasone treatment, 328
Complement C4, human:
  regulation of synthesis, 509
Complement regulatory activity, cell-surface bound:
  and tumour survival, 522
Cytokine production:
  by pig intestinal cells, 577
  in murine graft-versus-host reaction, 211
Cytomegalovirus, murine:
  induction of bone marrow atrophy, 410
Cytotoxic T lymphocyte, recognition of infected targets:
  cell-cycle dependence, 561
Cytotoxic T-lymphocyte epitope:
  naturally occurring, 337

Dendritic cell subsets:
  in human blood, 487
Dendritic cells, rat thymic:
  ontogeny of, 75

85C antigen on mycobacteria:
  and monocyte uptake of microbeads, 445
Eosinophils, priming:
  adhesion to fibronectin via $\alpha_4\beta_1$, 222
Experimental autoimmune uveoretinitis:
  suppression by intranasal administration of retinal antigens, 625

FcγRII activity, neutrophil:
  effect of proteolytic enzymes, 632
Foot-and-mouth disease virus (FMDV) peptides:
  induction of cross-reactive immunity, 1

Germinal centre cells:
  absence of intracellular antigen, 154
Glucocorticosteroid, topical:
  inhibition of cells expressing interleukin-4 mRNA in nasal mucosa, 192
Granulocyte–macrophage colony-stimulating factor:
  platelet-activating factor, 51
Granulocytes, human:
  respiratory syncytial virus antibody complexes on cytokine release and respiratory burst, 184
Granzyme A:
  expression in leishmaniasis, 255

Heat-shock protein 65, mycobacterial:
  expressed by transgenic antigen-presenting cell, 244
Hepatoma cells, human:
  accessory cell function, 215
Histoincompatiblity-associated differences:
  in allograft infiltrating T cells, 149
HLA class II molecules:
  on cells in human central nervous system, 535

ICAM-1 gene expression:
  in human hepatocytes, 571
IgA, human:
  role of carbohydrate in complement fixation, 321
IgE production, in *Nippostrongylus brasiliensis* infection:
  and suppressed interferon-γ, 427
IgE receptor, low affinity:
  on bronchial epithelial cells, 506
Interleukin-1 receptor, on astrocytes:
  regulation by cytokines and Theiler's virus, 178
Interleukin-2:
  chemoattractant activity for lymphocytes, 134
Interleukin-4 mRNA:
  expressed as T lymphocytes and mast cells in nasal mucosa, 200
Interleukin-6:
  role in protection during mycobacterial infection, 361
Interleukin-8 and RANTES:
  induction of basophil adhesion, 473
Interleukin-10:
  inhibition of motility in B lymphocytes, 376

Intraepithelial lymphocytes, murine:
heterogeneity, 611

Leucosialin:
characterization on human mast cell line HMC-1, 638
Lipids, dietary:
effect on lymphocytes, 603
Locomotor phenotypes:
of CD4 and CD8 lymphocytes in three-dimensional collagen lattices, 617
Lymphokine-activated killer (LAK) resistance:
of human cytomegalovirus-infected fibroblasts, 171
Lymphomyeloid cells:
in livers of *Plasmodium yoelii*-infected mice, 645

Macrophage priming/activation:
gene regulation, 42
Malaria, protection against:
induced by intrasplenic injection with infected hepatocytes, 164
Mannan-binding protein:
in amniotic fluid, 439
Mast cell secretion:
interactions of cytokines, 70
Mast cells:
natural suppressor activity, 482

Natural killer:
role in *Listeria* infection, 106
Natural killer cell activity:
decreased by tick salivary gland extracts, 113
Neutrophils, human:
priming of oxidative burst, 465
Nitric oxide:
and inflammation, 370
Nitric oxide synthase:
in adult rat thymus, 659
Nitrite production:
by macrophages, 457
NOD mice, thymus:
effect of transgenic expression, 275

Peptides and immunosuppressive effect of maternal antibodies, 16
Perforin:
susceptibility of peripheral blood leucocytes and cell lines to killing by, 555
Pulmonary surfactant:
inhibition of interferon activation of monocytes, 450

Respiratory syncytial virus:
infection of human pulmonary epithelial cell line, 126
Retrovirus:
induction of immunosuppression via blocking dendritic cell migration, 82

Self-peptides, major histocompatibility class II bound:
separation of, 529
Self-reactive T cells:
deletion prevented by anti-CD11a, 301
Substance P:
effect on interleukin-8 production by human polymorphonuclear leucocytes, 63
Superantigens:
cytokine production and cytotoxicity, 117
Suppressor macrophages:
role in inhibition of T-cell proliferation by poly I:C, 238
Suppressor T cells:
extrathymic activation by self-antigens, 365
Syphilis, congenital and neonatal, in guinea-pigs:
immune reponses, 404

$\gamma\delta$ T cells, ovine:
antigen recognition and activation, 229
T-cell activation, polyclonal:
by silicate, 332
T-cell peptides, bovine:
for three *Mycobacterium bovis* antigens, 9
T-cell receptor ($\alpha\beta$) gene usage:
and immunodominance, 343
Tick infestation, of mice:
immunological changes in skin, 596
Tolerance, intravenously induced:
and energy of antigen-specific T lymphocytes, 294
Transforming growth factor-β_1, in lung fibrosis, 584
Tumour-necrosis factor-α, role in inflammation:
dependence upon Th1/Th2 cytokine balance, 591

Ultraviolet B radiation:
induction of tolerance and impairment of contact hypersensitivity, 140

V gene analysis of anti-cardiolipin antibodies, 494
Vβ8 gene usage:
in response to TNP, 99